Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.


Volume 143 Issue 9 September 2010
ISSN: 0006-3207
ELSEVIER
BIOLOGICAL
CONSERVATION
Editor-in-Chief: Richard B. Primack
Editors: Richard Corlett, Markus Fischer, Andrew B. Gill,
Julie Lockwood, Jean P. Metzger, Daniel Oro,
Andrew S. Pullin, Navjot S. Sodhi

This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

http://www.elsevier.com/copyright

Author's personal copy

Biological Conservation 143 (2010) 2080–2091

Contents lists available at ScienceDirect

Biological Conservation

journal homepage: www.elsevier.com/locate/biocon

Woody species diversity in temperate Andean forests: The need for new conservation strategies

Adison Altamirano [a,*], Richard Field [b], Luis Cayuela [c], Paul Aplin [b], Antonio Lara [d], José María Rey-Benayas [e]

[a] Departamento de Ciencias Forestales, Universidad de La Frontera, P.O. Box 54-D, Temuco, Chile
[b] School of Geography, University of Nottingham, Nottingham NG7 2RD, United Kingdom
[c] Departamento de Ecología, Centro Andaluz de Medio Ambiente, Universidad de Granada – Junta de Andalucía, Granada 18006, Spain
[d] Instituto de Silvicultura, Universidad Austral de Chile, P.O. Box 567, Valdivia, Chile
[e] Departamento de Ecología, Edificio de Ciencias, Universidad de Alcalá, 28871 Alcalá de Henares, Spain

ARTICLE INFO

Article history:
Received 6 April 2009
Received in revised form 20 April 2010
Accepted 23 May 2010
Available online 16 June 2010

Keywords:
Biodiversity
Hotspot
Natural protected areas
Species richness
Spatial modelling

ABSTRACT

Chile has more than half of the temperate forests in the southern hemisphere. These have been included among the most threatened eco-regions in the world, because of the high degree of endemism and presence of monotypic genera. In this study, we develop empirical models to investigate present and future spatial patterns of woody species richness in temperate forests in south-central Chile. Our aims are both to increase understanding of species richness patterns in such forests and to develop recommendations for forest conservation strategies. Our data were obtained at multiple spatial scales, including field sampling, climate, elevation and topography data, and land-cover and spectrally derived variables from satellite sensor imagery. Climatic and land-cover variables most effectively accounted for tree species richness variability, while only weak relationships were found between explanatory variables and shrub species richness. The best models were used to obtain prediction maps of tree species richness for 2050, using data from the Hadley Centre's HadCM3 model. Current protected areas are located far from the areas of highest tree conservation value and our models suggest this trend will continue. We therefore suggest that current conservation strategies are insufficient, a trend likely to be repeated across many other areas. We propose the current network of protected areas should be increased, prioritizing sites of both current and future importance to increase the effectiveness of the national protected areas system. In this way, target sites for conservation can also be chosen to bring other benefits, such as improved water supply to populated areas.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

Loss of biodiversity is one of the most serious environmental problems today because of the associated economic, scientific, amenity and ecosystem service losses and the irreversible nature of global extinction (Newton, 2007). Threats to biodiversity remain strong, in large part because of continued increase in the rate of human-mediated destruction and conversion of habitats (May et al., 1995; Nagendra, 2001; Newton, 2007). The need to preserve biodiversity is therefore urgent. One of the main actions to protect biodiversity is to create or expand protected areas (Murphy, 1990; Nagendra, 2001). Selection of areas for conservation should take into consideration the representation and persistence of key attributes within sets of areas (Araújo, 1999). Species diversity is often used as a target attribute of biological communities to determine areas of high conservation value (De Vries et al., 1999; Luoto et al., 2002; Armenteras et al., 2006; Cayuela et al., 2006a); although it is only one of the important variables, it often correlates with other key measures. In turn, species richness (by which we mean the number of species in a given area), which is both the simplest and most easily interpreted measure of species diversity, tends to correlate strongly with the other measures (Whittaker et al., 2001). Explaining patterns of species richness is, however, a complex challenge because the diversity results from many interacting factors that operate at different spatial and temporal scales (Diamond, 1988; Willis and Whittaker, 2002).

At fine scales, a variety of variables typically account for (or at least correlate with) spatial diversity patterns (Whittaker et al., 2001; Field et al., 2009). These fine-scale correlations are usually weaker than those at broad scales (Field et al., 2009). Changes in elevation, slope or exposure can determine the ecological response of individual species and therefore contribute to overall changes in species richness (Luoto et al., 2002). Human activities also influence the shape of geographical patterns of diversity in intensively

* Corresponding author. Tel.: +56 45 734159; fax: +56 45 325634.
E-mail addresses: aaltamirano@ufro.cl (A. Altamirano), Richard.Field@nottingham.ac.uk (R. Field), lcayuela@ugr.es (L. Cayuela), Paul.Aplin@nottingham.ac.uk (P. Aplin), antoniolara@uach.cl (A. Lara), josem.rey@uah.es (J.M. Rey-Benayas).

0006-3207/$ - see front matter © 2010 Elsevier Ltd. All rights reserved.
doi:10.1016/j.biocon.2010.05.016

Author's personal copy

managed regions (Lawton et al., 1998; Ramírez-Marcial et al., 2001; Cayuela et al., 2006b; Hall et al., 2009).

At broader spatial scales, patterns of species richness are correlated strongly with climatic variables (Currie, 1991; O'Brien, 1998; O'Brien et al., 2000; González-Espinosa et al., 2004; Field et al., 2005). If climate directly or indirectly determines patterns of richness, then when the climatic variables change, richness should change in the manner that spatial correlations between richness and climate would predict (Acevedo and Currie, 2003; Venevsky and Veneskaia, 2003; Field et al., 2005). This might have important consequences for long-term conservation, since prioritization of highly diverse habitats today might not be effective in preserving future hotspots of species richness in the face of climate change.

In this study, we develop empirical models to investigate present and future spatial patterns of woody species richness in temperate forests in south-central Chile. We follow the lead of Cayuela et al. (2006a), who developed a predictive model using a similar approach, which allowed identification of high-priority areas for conservation of tropical forests in areas where the accessibility was limited. Our models include information obtained at multiple spatial scales, including field sampling, climate, topography and land-cover variables. The applied goals of this research are to inform attempts to prioritize the extant forest patches in the region and to provide recommendations for their conservation. This is of paramount importance as these forests are included in the Global 200 initiative launched by the World Wildlife Fund and the World Bank (Dinerstein et al., 1995), which focuses on the most threatened eco-regions in the world. In addition, these forests have been classified as one of the world's biological hotspots, e.g. by Myers et al. (2000), because of their high degree of endemism and presence of monotypic genera (Arroyo et al., 1996; Smith-Ramírez, 2004). The temperate forests of Chile are specifically considered to be vulnerable to impacts of climate change (IPCC, 2001; Pezoa, 2003). Paradoxically, in Chile, at broad scales the amount of land dedicated to conservation is inversely correlated with the number of species and endemism (Armesto et al., 1998). Thus, more than 90% of the 14 million hectares of protected land (CONAF et al., 1999) is concentrated in high latitudes (>43°), leaving unprotected a large proportion of high-biodiversity areas (Armesto et al., 1998). Here, we investigate whether the inverse relationship between amount of conserved land and numbers of species is true at a smaller spatial scale. For all these reasons, establishing guidelines for prioritization of natural protected areas is a crucial step towards biodiversity conservation in this important eco-region.

The specific objectives of this study are: (a) to assess the independent and joint contribution of different groups of variables in describing the variation in woody species richness in the study area, thereby increasing our knowledge and understanding of Chile's temperate Andean forests; (b) to develop a model to estimate present-day, fine-scale woody species richness across the study area; (c) to develop a model to predict the effects of climate change on woody species richness; and (d) to use the models to evaluate the effectiveness of the currently protected areas for maintaining biodiversity both now and in the face of climate change. The models we develop can also be used to inform future modification of the protected area network and to facilitate forest restoration programmes.

2. Materials and methods

2.1. Study area

Our study was conducted in the Maule region of Chile, which lies mainly in the Andean area between 35° and 36° latitude south (Fig. 1). The study area covers approximately 270,000 ha and is between 200 and 3900 m.a.s.l. The predominant soils are volcanic in origin, with different degrees of development (Schlatter et al., 1997). The predominant climate is of the Mediterranean type, with annual precipitation averaging between 700 and 1300 mm and concentrated mostly during the winter season, and an average annual temperature of 9 °C (Pezoa, 2003).

The area is characterized by the presence of secondary and old-growth forests (dominated by species like *Nothofagus obliqua*, *Nothofagus glauca*, *Nothofagus dombeyi* and sclerophyllous species over 2 m high and >50% coverage), shrublands (composed mainly of low-height sclerophyllous species such as *Criptocarya alba*, *Quillaja saponaria* and *Lithraea caustica*), exotic plantations (mainly of *Pinus radiata*), agricultural lands, herbaceous vegetation, grasslands, and other types of land-cover such as bare land, urban areas and water bodies (Appendix A) (CONAF et al., 1999; Altamirano et al., 2007). The intensification of land use, particularly firewood extraction and selective logging, has caused much deforestation and forest disturbance, which may have a negative impact on biodiversity (Lara et al., 1996, 2003; Olivares, 1999; Echeverría et al., 2006).

The national protected areas system of Chile comprises 96 sites, totalling approximately 14 million hectares and representing 19% of the land (CONAF et al., 1999). The three main types of protected area are National Parks, National Reserves and Natural Monuments. National Reserves are medium-sized areas that are protected with the aim of conserving species, soils and hydrological resources; sustainable natural resource use is allowed. There are two of these reserves in our study area: Altos de Lircay (approximately 12,000 ha) in the north, and Los Bellotos (approximately 400 ha) in the south (Fig. 1).

2.2. Field sampling and estimation of woody diversity

The study area was divided into approximately 700 cells, each 2×2 km. Of these, 82 were selected via a random sampling scheme stratified by vegetation structure (see below), to contain field plots. One field plot was located in each of these 82 cells so that it was as close to the centre of the plot as possible, given the constraints that it was within the most representative vegetation structure in terms of percentage cover inside the cells, and was accessible. The plots provided good coverage of the main vegetation and soil types, and of the elevational range. In a pilot study, the numbers of species in 10 circular plots of 500 m^2 and 250 m^2 were compared. No significant differences were found (Student's paired t-test, $t = 2.3$, $P = 0.16$), so in order to allow greater replication, 250 m^2 (i.e. 9 m radius) was set as the plot size. This also better matched the resolution of the ASTER imagery used (see next section). The 82 plots were sampled in 2005 and 2006. In each, all trees and shrubs with a height greater than 1.4 m were identified to species (see Appendix A), counted and measured; from this, we calculated basal area. Fisher's alpha index, Shannon's diversity index and species richness (number of species observed) were calculated for each sample. Fisher's alpha and Shannon's indices were, however, highly correlated with species richness ($r = 0.92$, $P < 0.0001$; $r = 0.87$, $P < 0.0001$ respectively). Because of this strong similarity and the ease of interpretability, we only report results for species richness.

2.3. Explanatory variables

To model species richness we focused on six climatic variables, two topographic variables and three land-cover variables (Table 1). We initially obtained 19 climatic variables from the WorldClim database (www.worldclim.org). WorldClim is a set of global climate layers (climate grids) with a spatial resolution of 1×1 km (Hijmans et al., 2005). This set includes 19 temperature, rainfall and bioclimatic variables. The bioclimatic variables were derived from the monthly temperature and rainfall values in order to be

Author's personal copy



Fig. 1. Map of the study area in the Andean range.

more biologically meaningful, and represent annual trends in seasonality and extreme or limiting environmental factors (Hijmans et al., 2005). We carefully examined the correlation matrix to determine the degree of collinearity and redundancy between these climatic variables (and the other explanatory variables), as well as their correlations with species richness. We additionally performed a hierarchical cluster analysis of these variables in order to identify groupings of correlated explanatory variables. To achieve this, we used the 'Hmisc' library (Harrell et al., 2009) of the R environment (R Development Core Team, 2009), defining a threshold of Spearman's $\rho = 0.6$. We combined this information with theoretical considerations to select climatic variables for further analysis that would minimise multicollinearity, while being expected to account best for species richness, as recommended by Carsten F. Dormann (pers. comm.). Multicollinearity tends both to promote statistical artefacts (resulting in false model accuracy) and to cause unstable parameter estimates, which are particular problems when making predictions of future diversity. Thus we chose the following climatic variables for the regression analyses (Table 1): minimum temperature of the coldest month (T_{min}), temperature seasonality (T_{seas}), mean annual precipitation (P_{an}), mean precipitation of the driest month (P_{min}) and precipitation seasonal-

Table 1
Climatic, topographic and land-cover variables used to model the spatial variation in woody species richness in the study area. Values given are for the 82 field plots. (Code = abbreviation used, St. dev. = standard deviation, *r* (WSR) = Pearson's correlation coefficient for the relationship with woody species richness, *r* (TSR) = correlation with tree species richness, *r* (SSR) = correlation with shrub species richness, CV = coefficient of variation.)

Variable and unit of measurement	Code	Mean	St. dev.	Min.	Max.	*r* (WSR)	*r* (TSR)	*r* (SSR)
Response variables								
Woody species richness	n.a.	8.3	3.1	2	16	n.a.	0.83***	0.63***
Tree species richness	n.a.	5.0	2.4	0	11	0.83***	n.a.	0.10 n.s.
Shrub species richness	n.a.	3.3	1.7	0	8	0.63***	0.10 n.s.	n.a.
Climatic variables								
Elevation (m) – untransformed	ELEV	674	292	288	1603	−0.50***	−0.63***	−0.03 n.s.
Minimum temperature of coldest month (°C)	T_{min}	0.31	1.29	−3.0	2.2	0.50***	0.60***	0.05 n.s.
Temperature seasonality (st. dev.)	T_{seas}	4.54	0.09	4.39	4.74	−0.28*	−0.42***	0.09 n.s.
Annual precipitation (mm)	P_{an}	1080	76	853	1260	0.22*	0.40***	−0.16 n.s.
Precipitation of driest month (mm)	P_{min}	15.8	1.6	11	18	−0.47***	−0.53***	−0.11 n.s.
Precipitation seasonality (CV × 100)	P_{seas}	83.9	1.9	78	88	0.33**	0.44***	−0.02 n.s.
Topographic variables								
Aspect (degrees from north)	ASPE	83.2	55.7	1	180	0.02 n.s.	0.17 n.s.	−0.21 n.s.
Slope (degrees)	SLOP	14.2	8.5	1	38	−0.18 n.s.	−0.11 n.s.	−0.17 n.s.
Land-cover variables								
Normalised difference vegetation index	NDVI	0.678	0.196	0.056	1.000	−0.13 n.s.	0.01 n.s.	−0.26*
Normalised difference infrared index	NDII	0.719	0.064	0.455	0.825	−0.23*	−0.11 n.s.	−0.26*
Vegetation structure[a]	VST	n.a	n.a.	1	4	n.a.	n.a.	n.a.

n.a. = correlation is not applicable.
n.s. = not significant.
[a] Categorical data: four categories (1 = open shrubland, 2 = dense shrubland, 3 = arborescent shrubland, 4 = forest).
* $P < 0.05$.
** $P < 0.01$.
*** $P < 0.001$.

Author's personal copy

ity (P_{seas}). Mean annual temperature was strongly correlated with T_{min} ($r = 0.98$); we chose T_{min} because it is very similar to minimum monthly potential evapotranspiration calculated by the Thornthwaite method, which previous empirical and theoretical work has shown to be a good predictor of woody species richness (e.g. Field et al., 2005). In our dataset, T_{min} correlated more strongly with species richness than T_{mean}, supporting our reasoning. P_{min} is appropriate in climates where precipitation is lowest during the summer months, as in our study area, because it represents a strong constraint on growth.

Elevation was derived from a digital elevation model, with a spatial resolution of 90 × 90 m, based on the Shuttle Radar Topography Mission (SRTM). The SRTM data are available from the Global Land Cover Facility (GLCF) website (http://www.landcover.org). It was classed as a climate proxy for several reasons. Elevation is a powerful and very precise determinant of small-scale climatic variation, particularly temperature; this is because of the close association between temperature and elevation that results from the effects of the adiabatic lapse rate. The relationship between elevation and precipitation is less strong, more indirect and more complex. In this study, elevation, with its resolution of 90 × 90 m, is much more precisely measured than the WorldClim variables (resolution 1 × 1 km), so it can be expected to model climate (particularly temperature) well for the field plots. Given the scale of the field plots and the nature of the study area, elevation was also a poor topographic measure, indicating nothing about topographic heterogeneity, nor about aspect. This reasoning is backed up by the fact that, in our dataset, elevation was not correlated with topographic variables ($r = 0.035$ and -0.044 for slope and aspect respectively), but was almost perfectly inversely correlated with T_{min} and mean annual temperature ($r = -0.95$ and -0.94) despite the difference in resolution. The correlation between elevation and precipitation was moderate ($r = -0.55$ for P_{an}; $r = -0.71$ between 1/elevation and P_{min}).

The topographic variables used were aspect and slope (Table 1). Aspect was measured as degrees from north. These variables were derived from the digital elevation model.

We performed image analysis on remotely sensed imagery, acquired in March 2003 by the Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER), to identify different forest types in relation to their degree of disturbance and to calculate different vegetation indices. We georeferenced the image using 43 control points derived from vector maps of roads and rivers, obtained from the Native Vegetation Survey (CONAF et al., 1999), resulting in an estimated error of less than one pixel. We atmospherically corrected the image using the dark pixel subtraction method and features such as water bodies (Mather, 1999).

We performed a supervised land-cover classification (Aplin, 2004) of the image using the maximum likelihood algorithm (Lillesand et al., 2004). Four different types of vegetation structure (VST) were identified in relation to human disturbance: open shrubland (VST1), dense shrubland (VST2), arborescent shrubland (VST3) and forest (VST4), which includes old-growth forest, secondary forest and an intermediate condition (Altamirano and Lara, 2010). Other land-cover types were excluded from the analyses and predictions reported herein. Training sites were selected using different sources of information such as vegetation maps, aerial photographs from 2003 and field visits conducted between 2004 and 2006. The overall accuracy of the supervised classification was 92%. The lowest accuracy was obtained for shrubland and secondary forests; this was because of spectral confusion between these two classes. The pre-processing and classification of remotely sensed data were performed using the ERDAS Imagine 8.4® software (ERDAS, 1999).

In addition, we calculated two spectral indices: the normalised difference vegetation index (NDVI) and the normalised difference infrared index (NDII). The NDVI was calculated as the difference between the near-infrared and red reflectances divided by their sum, which represents a measure of vegetation productivity (Turner et al., 2003; Aplin, 2005). The NDII was calculated as the difference between the near-infrared and mid-infrared reflectances divided by their sum, which is related to the hydric stress (Bannari et al., 1995; Gao, 1996). We did not use spectral bands from the ASTER image because they were both highly correlated with, and less interpretable than, NDVI and NDII ($|r| > 0.85$, $P < 0.001$).

2.4. Statistical analyses

To inform subsequent analysis, we used variance partitioning to explore the independent and joint contribution of all available explanatory variables, including all 19 WorldClim variables and elevation ('climatic' category), the spectral bands from the ASTER image and derived land-cover variables ('land-cover') and slope and aspect ('topographic') in accounting for spatial variation in woody species richness. The partition of the variance is derived from partial redundancy analyses (RDA) and was used to determine the proportions that could be attributed to the single and combined effects of explanatory variables (Legendre and Legendre, 1998), using adjusted R^2 ratios (Peres-Neto et al., 2006). These analyses were computed using the 'vegan' library (Oksanen et al., 2008) of the R environment (R Development Core Team, 2009).

The results of the variance partitioning informed the selection of variables for modelling, in which we used multiple regression to develop models to predict woody species richness in the study area. Most regression analysis was performed with the S-PLUS 6.0 software (Insightful Corporation, 2001); spatial analysis was conducted using SAM (Rangel et al., 2006). Before performing multiple regression, we examined the correlation matrix and the hierarchical cluster analysis. We noted explanatory variables that were highly correlated ($|r| > 0.6$) and the clusters of such correlated variables, which might therefore lead to problems associated with multicollinearity. We used the same mix of theoretical and statistical considerations as described above for selection of WorldClim variables, to determine which of all the explanatory variables could be combined in any one model. We also calculated the variance inflation factor (VIF) for all terms in all multiple-regression models, to quantify any remaining multicollinearity, using a maximum allowable level of VIF of 4. In addition, we examined the correlations between species richness and all selected explanatory variables (Table 1). Further, because relationships between species richness and environmental variables are often curvilinear (Austin, 1980) and interactive (Francis and Currie, 2003), we included quadratic and cubic terms in the models, as well as some interactions expected from previous research (e.g. between temperature and water variables).

Our modelling procedure was step-wise and manual (Murtaugh, 2009), using a combination of model building and model simplification. We produced the first model by building from the null model (the mean), adding terms in order of explanatory power, defined as the change in residual sum of squares resulting from the addition of individual terms to the current model. Model building finished when no more terms were both significant and reduced AIC (Akaike Information Criterion) (Venables and Ripley, 2002; Anderson and Burnham, 1999). We then used a similar procedure, but with different starting variables (chosen according to variable type and variance accounted for) and different orders of addition. We also produced a series of models by simplifying from various maximal models (Crawley, 2002). It was necessary to simplify from more than one maximal model because the sample size of 82 did not support highly complex models. We compared all models obtained statistically using P-values, AIC and the proportion of variation accounted for (R^2). We also used 'Model Selection

Author's personal copy

and Multi-model Inference' in SAM to rank over 16,000 possible models by AICc and AIC-weights and to calculate the 'importance' of each variable across all the models. This combination of model fitting approaches allowed confidence in the robustness of the results. Selection of the 'best' models was based on both theoretical criteria (plausibility, generality, simplicity, parsimony) and statistical strength (O'Brien et al., 2000).

We used Moran's I to evaluate the spatial autocorrelation of the residuals of the fitted models. Finding no residual spatial autocorrelation means that we can assume the significance values to be reliable, and that we do not need to introduce the further uncertainties (coefficient instability) associated with spatial regression (Bini et al., 2009). We checked the model residuals for normality using histograms and the Kolmogorov–Smirnov test. We assessed homoscedasticity via residual plots, and we mapped model residuals to examine their spatial patterning. To validate the predictive power of the models, we used a bootstrap approach for each model. This method generates new samples with replacement from the original sample, allowing a quantification of the error introduced by data uncertainty as well as model estimation procedure (Quinn and Keough, 2003).

We used the resulting regression models to predict current species richness values for parts of the study area where there were no field plots. This was done for every pixel in the ASTER imagery and involved using the coefficients from the models and substituting the applicable values for the explanatory variables, to calculate predicted current species richness.

2.5. Climate-change scenario

We used the coefficients derived from modelling current woody species richness to predict future species richness across the study area, using climatic data obtained for a climate-change scenario. Climate scenarios are guesses of future climates, based on assumptions about future emissions of greenhouse gases and other pollutants, and obtained via general circulation models, such as CCCMA, HadCM3 and CSIRO. We used projected climate data for 2050, from the Hadley Centre's climate model (HadCM3 Worldclim implementation) under the low (B2a) CO_2 emissions scenario (Zhang and Nearing, 2005), obtained from WorldClim. We used scenario B2a because it emphasizes more regionalized solutions to economic, social, and environmental sustainability (Zhang and Nearing, 2005).

2.6. Conservation value

To analyse conservation value we produced maps categorizing the predicted current woody species richness into three levels: high (>8 species), medium (5–8 species) and low (<5 species). When examining tree species richness, we used >6, 4–6 and <4 species respectively. This was done for all pixels of the ASTER image (with a spatial resolution of 15 m) that had land-cover in one of the categories VST1, VST2, VST3 and VST4. Pixels classified as other categories were excluded from further consideration. Predicted future species richness was categorized using the same criteria and results were compared in terms of: (1) forest area occupied by each conservation value category now and in 2050; (2) overall forest area that will change to a different conservation value category by 2050; and (3) forest area in current natural protected areas assigned to different conservation value categories now and in 2050.

3. Results

We recorded 67 woody species (28 trees and 39 shrubs) in the field plots (Appendix A), with a mean (±SD) number of species per plot of 8.3 (±3.1), ranging from 2 to 16. We found significantly lower mean tree and overall (tree + shrub) species richness in VST1 (open shrubland) than in the other three categories (ANOVA, $P = 0.006$ and 0.017 respectively), but no differences between VST2 (dense shrubland), VST3 (arborescent shrubland) and VST4 (forest). Therefore for regression modelling we re-categorized the forest structure variable into two categories: VST1 and closed canopy (VST2, VST3 and VST4 combined) because this is more robust and parsimonious (Crawley, 2002). Interestingly, there was no significant difference between VST categories in either tree or shrub abundance.

The strongest single-variable correlates of both tree and overall species richness in the 82 field plots were T_{min}, P_{min} and ELEV (Table 1). None of the explanatory variables in Table 1 correlated with shrub species richness at the 1% significance level. At the 5% level only NDVI and NDII were significant, both correlating weakly and negatively with shrub species richness (Table 1); this effect was driven by the open shrubland ($r = -0.50$ for both NDVI and NDII) and was not significant for the denser woody vegetation categories. The difference in shrub diversity between the different VST categories was also not significant. Basal area of woody plants correlated negatively with shrub species richness ($r = -0.31$, $P = 0.005$) and band 5 of the ASTER image correlated positively ($r = 0.36$, $P = 0.0009$). Band 5 and basal area represented the strongest statistical model, with neither NDVI nor NDII significantly improving it, but this model only accounted for 18% of the variation, had non-normal residuals and contained potential circularity. Overall, then, we were unable to produce a satisfactory model of shrub species richness, which also did not correlate significantly with tree species richness (Table 1). Models of overall woody species richness were all qualitatively identical to, but quantitatively weaker than, those for tree species richness; they were driven by the tree species richness pattern, with shrub species richness effectively adding noise. We therefore focus on reporting the results for tree species richness.

Minimum temperature (T_{min}) correlated positively and ELEV negatively with tree species richness, both consistent with greater energy allowing more species. The correlation between tree species richness and P_{min}, however, was negative, both singly and when included in multiple-regression models. Both log and inverse transformations of ELEV improved the linearity of its association with tree species richness, 1/ELEV the more so, which also improved the normality of regression residuals compared with models using ln(ELEV). Using 1/ELEV made the relationship with tree species richness positive and increased the strength of the bivariate correlation to $r = 0.66$.

In variance partitioning, topographic, climatic, and land-cover variables accounted for 2%, 52% and 7%, respectively, of the adjusted variance of woody species richness (Fig. 2). Overlap between the categories in variance accounted for was minimal (Fig. 2), supporting our contention that elevation acts as a climatic, not topographic, variable in our dataset.

3.1. Predictive models of species richness

T_{min} correlated strongly with 1/ELEV ($r = 0.91$), so only one of the two variables could be used in the same regression model. We developed models independently using both variables. In all cases, as with simple correlation, 1/ELEV gave a closer fit with woody species richness (Model 1, Table 2a). However, for predicting species richness for the year 2050 we preferred models featuring T_{min} instead of 1/ELEV (Models 2a and b). Temperature has a direct physiological effect on species performance, while elevation is a surrogate variable for a mixture of influences, but driven by temperature (Guisan and Zimmermann, 2000; Pausas and Austin, 2001), so temperature is preferred on theoretical grounds. Most

Author's personal copy



Fig. 2. Venn diagram of the partition of the variation of tree species richness for climatic, topographic and land-cover variables. Table 1 shows which of the key variables were included in each category; other WorldClim data were included as 'climatic' and other variables derived from ASTER imagery were included as 'land-cover'. The rectangle represents the total variance of tree species richness while each circle represents a given group of explanatory variables. The adjusted R^2 (expressed as % of the variance in tree species richness) is presented for each part of the Venn diagram; if missing this fraction does not differ significantly from 0. Intersections between circles represent the fraction of the variance of tree species richness jointly accounted for.

of the effect of elevation is related to temperature and precipitation, and with climatic changes over the next 40 years, the regression coefficients derived from current conditions for elevation are not applicable to prediction for 2050. Therefore for prediction of future species richness, and for comparison of the conservation value of protected areas now and in the future, we used the best models that were based on T_{min} (Models 2a and 2b; Table 2b and c). VST was considered appropriate for 2050 because its main determinant is human activity (disturbance); its inclusion in Models 2a and b assumes no change in the disturbance regime during the first half of the 21st century.

The 'best' regression models (Models 1 and 2a,b) were selected on the grounds of theoretical plausibility, simplicity and statistical strength (secondary to the other two). These best models included two alternative models based on T_{min}: Models 2a and 2b (Table 2). These were statistically indistinguishable and both were ecologically plausible. The strongest effects are the same in both models, and in a reduced model with only VST and T_{min}. The first is a strong increase in tree species richness with increased T_{min}, of approximately 1 species per 1 °C. The second is approximately 1.5 fewer species in open shrubland than the other vegetation types. Thus the core of the models is the same; they differ in the final variable included, which in each case only accounts for an additional 4% of the variance (approximately). In Model 2a, this is T_{seas}, with a decrease of approximately 6 tree species for every 1 °C increase in seasonality (measured as the standard deviation; Table 1); this is ecologically plausible (Jocqué et al., 2010). In Model 2b, the third variable is P_{min}, with a decrease of approximately 1 tree species for every 3 mm increase in driest-month precipitation. Exploring this negative effect further (see also Table 1), we found a negative correlation ($r = -0.40$) between P_{min} and overall tree abundance, suggesting a competition or crowding effect, coupled with a more individuals effect (positive correlation, $r = 0.54$, between tree abundance and tree species richness). However, using data for basal area and average tree diameter for all plots in the dataset, we found no correlation between either variable and P_{min}. Nor did either basal area or average tree diameter correlate with tree species richness. So, while P_{min} could be measuring a competition effect, we are far from certain that it does indeed do so, or whether it is measuring another biologically meaningful effect such as inhibition of seed germination or seedling survival (Donoso, 1994), or whether the apparent effect is due to correlation with other important biological influences. Because P_{min} and T_{seas} are positively correlated ($r = 0.58$) and neither is even close to significant when the other is in the model, Models 2a and 2b are straight alternatives and we are unable satisfactorily to reject one in favour of the other. Therefore our predictive modelling was based on average predictions from the two models, hereafter referred to collectively as Model 2. This averaging of predictions, a form of ensemble forecasting (Araújo and New, 2007), should also increase the robustness of the predictions.

All models presented in Table 2 are statistically significant ($P < 0.0001$) and all rely on few explanatory variables, reducing the likelihood of artefact, which is particularly important when predicting future species richness. All the models met assumptions of homoscedasticity and normality of residuals. For all the models, Moran's I values for residuals were not significant for any of the short distance classes (Fig. 3), indicating no inflation of degrees of freedom resulting from spatial autocorrelation, and the absence

Table 2
'Best' models for predicting spatial variation in tree species richness in the study area: (a) Model 1 – model for predicting current species richness (best model using all available variables); (b) Model 2a – model for predicting future species richness and for comparison of predictions; (c) Model 2b – alternative model for predicting future species richness and for comparison of predictions. D.f. = degrees of freedom; VIF = variance inflation factor; R^2 = proportion of the variance accounted for (tested by deletion from the model); AIC = Akaike Information Criterion; root mean square error (RMSE) = square root of the error variance. All predictions are for plots of 250 m^2. See Table 1 for full variable names and units.

Model	Coefficient	D.f.	VIF	*t*-value	*p*	R^2	AIC
Null model		81					383.5
(a) Model 1 (RMSE: 3.0)							
Intercept	−0.34			−0.53	0.597		
1/ELEV(in km)	2.35	1	1.00	8.07	0.000	0.41	376.7
VST	1.59	1	1.00	3.33	0.001	0.07	338.2
Overall model		2			0.000	0.50	329.6
(b) Model 2a (RMSE: 3.3)							
Intercept	30.18			2.79	0.007		
VST	1.54	1	1.01	3.09	0.003	0.07	344.5
T_{min}	0.93	1	1.21	5.40	0.000	0.20	361.0
T_{seas}	−5.88	1	1.20	−2.47	0.017	0.04	341.2
Overall model		3			0.000	0.47	337.3
(c) Model 2b (RMSE: 3.3)							
Intercept	9.51			2.90	0.005		
VST	1.51	1	1.01	3.02	0.003	0.06	344.8
T_{min}	0.81	1	1.66	3.98	0.000	0.11	350.9
P_{min}	−0.38	1	1.65	−2.31	0.024	0.04	341.2
Overall model		3			0.000	0.46	338.0

Author's personal copy

of intrinsic autocorrelation that could disturb the Type I error rates and the coefficient estimates. In other words, the regression models have accounted for the spatial autocorrelation present in the species richness data. This also means that our predictive model is of the type considered the best for predicting responses to climate change by Algar et al. (2009): they concluded that the most accurate predictions of shifts in species diversity in response to climate change are obtained via the single best richness–environment regression model, after accounting for the effects of spatial autocorrelation. Further, our model has the advantage that the spatial autocorrelation is accounted for via ordinary least-squares regression, so that there is no chance of real effects being 'corrected for' while removing spatial autocorrelation in spatial regressions.

3.2. Predicted species richness

Present-day woody species richness was predicted for the whole study area using Model 1 (Fig. 4a) and Model 2 (Fig. 4b). Model 2 predicted slightly higher species richness on average than Model 1, but the spatial patterns were very similar. The areas of higher predicted richness at this scale (250 m^2) are concentrated mainly in the western locations of the study area and in valleys, at lower elevation and higher temperatures. These areas are dominated by shrubland and arborescent vegetation. The two protected areas in the study area have relatively low levels of predicted current species richness (Fig. 4a and b).



Fig. 3. Correlograms for tree species richness, fitted values and residuals of the 'best' regression models: (a) Model 1, (b) Model 2a, (c) Model 2b. Equal distance classes; only classes with $n > 100$ shown. See Table 2 for model specifications ($n = 82$ cells).

Our predictions for 2050 (Fig. 5a) suggest that the higher ground in the east of the study area will increase in tree species richness, while the lower ground in the west will decrease. Thus the species-richness gradient across the study area is expected to persist but weaken (compare Fig. 5a with Fig. 4b) with climate change. Overall, of the 1296 km^2 for which we made predictions, a net loss of species was predicted for 490 km^2 (38%) and a net gain for 698 km^2 (54%), the remainder staying approximately constant. Using our categories for conservation value, 58% of the pixels (each 225 m^2) were predicted by Model 2 to have present-day woody species richness in the low category (0–3 species), with 31% having more than 6 species (Fig. 6a). Of all the pixels, 34% were predicted to change from low to medium conservation value, while 16% were predicted to change from high to medium (Fig. 6a).

Only 29.0 km^2 of the land currently designated as protected areas is covered by woody vegetation, as judged by our analysis of the ASTER image. All of this area is currently in the low conservation priority (value) category, according to Model 2. Our map predictions suggest that 8.6 km^2 (30%) of the protected area will improve to the medium category by 2050, the rest remaining 'low' (Fig. 6b).

4. Discussion

We found that the highest tree species richness occurs in low and medium elevation areas, with the highest minimum temperatures, and where there is relatively dense woody vegetation cover. The protected areas within the study area contain very low tree species richness and our modelling suggests that the areas of highest tree conservation value are far from the currently protected areas. Our predictions for changed climate indicate reduced tree diversity where it is currently high and increased diversity where it is currently low. The currently protected areas may therefore slightly increase in tree conservation value over the next 40 years, but will still be relatively low in diversity. Meanwhile, the areas of greatest species richness are predicted to suffer losses, thereby degrading in conservation value. The resulting predominance of areas of relatively average conservation value suggests a need for the conservation of greater areas of forest. Greater connectivity of patches of woody vegetation may also be important. Although the protected areas may be important for species other than woody plants, their continued low value for tree species conservation is of great conservation concern because Chile has more than half of the temperate forests in the southern hemisphere (Donoso, 1994), because of the uniqueness of these forests (Smith-Ramírez, 2004), and because of the high levels of threat to these forests (Dinerstein et al., 1995).

These concerns about tree conservation that arise from our species richness modelling are backed up by our field observations of threatened species within our study plots. We recorded 4 threatened species, all of which are trees: *Nothofagus glauca*, *Austrocedrus chilensis*, *Beilschmiedia berteroana* and *Cytronella mucronata*. These species have a restricted distribution and highly specific habitats (Hechenleitner et al., 2005). Migration capabilities for these species under climate change may well be limited. *N. glauca* (by far the most common of the four in our field plots) is restricted largely to the Maule region and is found mainly in intermediate elevation sites (Hechenleitner et al., 2005), so may not be much affected by climate change. However, *B. berteroana* may be negatively affected by climate change because its habitat is coincident with sites where species richness is expected to decrease. To aggra-

Author's personal copy



Fig. 4. Map of predicted current woody species richness in 225 m^2 pixels. (A) Current tree species richness according to Model 1; (B) current tree species richness according to Model 2 (average of predictions from Models 2a and 2b); (C) current conservation priority (value) category, as defined by tree species richness predicted by Model 2 (low = <4, medium = 4–6, high = >6). No colour means no prediction because the pixel is not currently classed as any of the land-cover types in our analyses. See Table 2 for model specifications.

vate the problem, only 8 sub-populations of this species have been identified in the country (Hechenleitner et al., 2005). The other two threatened species have wider distributions and may therefore be less vulnerable to climate change. An additional consideration is that many mountain plants reproduce vegetatively and grow slowly; consequently they are likely to take a long time to disperse into new, climatically suitable areas (Trivedi et al., 2008).

There are continuing threats to temperate Andean forests and their biodiversity, such as hydroelectric-power projects and the rapid growth of the exotic plantations industry (Lara et al., 2003). In recent years, exotic plantations have expanded specifically in the south-central temperate forests of Chile (Echeverría et al., 2006) because of the growth of the pulp and wood industry (Lara et al., 2003). These developments may facilitate the establishment and

Author's personal copy



Fig. 5. Map of predicted tree species richness in 2050, in 225 m^2 pixels. (A) Tree species richness in 2050 according to Model 2 (average of predictions from Models 2a and 2b); (B) change in tree species richness from now to 2050, according to Model 2; (C) uncertainty for 2050 tree species richness predictions (absolute difference between the predictions of Models 2a and 2b); (D) conservation priority (value) category in 2050, as defined by species richness predicted by Model 2 (low = <4, medium = 4–6, high = >6). No colour means no prediction because the pixel is not currently classed as any of the land-cover types in our analyses. See Table 2 for model specifications.



Fig. 6. Current forest area by conservation priority (value) category (horizontal axis labels), and how these conservation priorities will change in the year 2050 (shading), according to Model 2 (average of predictions from Models 2a and 2b). (A) In the study area and (B) in the current protected areas. See Table 2 for model specifications.

invasion of alien species, which may also be enhanced by predicted rises in the frequency of natural disturbances (e.g. forest fires), and ultimately reduce the cover of native vegetation (Pickering et al., 2008). Studies in Chile have shown that alien species are moving into native forests in national parks in mountain areas (Pauchard and Alaback, 2004). Given these various forms of disturbance in the study area, our results suggest that protected areas are important for conservation: we found that the most disturbed areas of woody vegetation have the lowest tree species richness, with no accompanying increase in shrub species richness. This suggests that one way of improving conservation is to minimize disturbance.

Our predictions, by necessity, assumed no change in protection/disturbance regime (land-cover type). Nonetheless, our coefficients for the disturbance variable (VST) can be used to explore future scenarios in which disturbance regimes do change in prescribed ways. Our assumption of no increase in disturbance may be optimistic, unless the protected area network is modified, or unless parts of the landscape not in protected areas are managed for woody plant conservation. We consider that both strategies should be implemented. New protected areas should be created, and because our prediction maps indicate that current high-priority sites are coincident with high-priority sites in 2050, we suggest that sites that have high tree species richness now should be targeted for national protection. In the study area, these sites include river valleys, and so this should help to ensure reliable supplies of clean water downstream. Such targeting is important: Babcock et al. (1997) demonstrated that enrolling land into a conservation programme on the basis of the lowest cost of purchasing land (as has been the case for many of Chile's protected areas) is a far less efficient use of taxpayers' money than targeting land on the basis of the cost–benefit ratio of that land. The application of newer approaches to protected area design could help stakeholders find designs that simultaneously maximize ecological, societal and industrial goals (Gonzales et al., 2003). Planning tools such as Sites

Author's personal copy

(Davis et al., 1999) and Marxan (Game and Grantham, 2008) represent good examples. Of high relevance to areas not formally protected, in 2008 Chile passed a new law that supports native forest management and biodiversity conservation. This law gives economic incentives to landowners to engage in biodiversity conservation. Again, the law can target high-priority conservation sites, as indicated by our prediction maps, for improved effectiveness (Macmillan et al., 1998). Subsidies to encourage landowners to manage their land in ways that increase the provision of non-market benefits may also be appropriate (Van der Horst, 2007).

Our research represents a starting-point, but more work is needed to inform conservation in the temperate Andean forests. First, our model should be seen as a tool, for addressing urgent conservation issues, that should be assessed, discussed and evaluated further. Also, we have not investigated individual species' requirements; for example, species-specific conservation measures for endemic and threatened species, including ex situ conservation, may be required. Woody species may have differential abilities to cope with climate change (Parolo and Rossi, 2008), and habitat connectivity may be important in enabling some to migrate. Species usually differ in their habitat requirements and habitat mosaics may be appropriate in meeting each species' needs (Drechsler et al., 2007). In this context, important future challenges for biodiversity conservation research are to investigate beta diversity and determine how much habitat heterogeneity is needed to maintain species diversity at coarser scales than in our study. Furthermore, we used climatic and topographic data that are widely used for this sort of analysis (WorldClim and SRTM). However, some environmental data sets may be less useful in some areas (i.e. rugged, remote and steep terrain) and scales (Peterson and Nakzawa, 2008). Therefore, further corroboration and testing of other source information will be necessary.

Our study adds to knowledge and understanding of species richness patterns and their correlates. Tree species richness correlated most strongly with temperature-related variables (elevation and minimum temperature), which is common at broad scales but less common at the finer scale of our study (Field et al., 2009). This may be because we sampled quite a large altitudinal range, and fits with the findings of Bhattarai and Vetaas (2003). The closer match, in terms of scale of measurement, between elevation and species richness, compared with climatic variables, probably explains the stronger correlation of species richness with elevation. The relatively small amount of variation accounted for by P_{min} and T_{seas} is probably due, in large part, to the fact that both vary little in the data for our study plots (Table 1). Despite their coarse scale of measurement, climatic variables performed well in accounting for tree species richness patterns, relative to the fine-resolution variables such as slope, aspect and NDVI. This supports the contention (Cayuela et al., 2006a) that broad-scale patterns (e.g. Hawkins et al., 2003; Field et al., 2005) can be replicated across altitudinal gradients at finer spatial scales. In addition, there was a positive correlation ($r = 0.54$) between tree abundance and tree species richness in our field plots; adding tree abundance to any of the final tree species richness models led to about a 5% increase in variation accounted for. This suggests a 'more individuals' effect, whereby more individuals tend to be associated with more species (Srivastava and Lawton, 1998; Currie et al., 2004). However, this was of little use for modelling because our best model of tree abundance contained only T_{min} and only accounted for 26% of the variation.

Our best tree species richness model accounts for approximately 50% of the variance, which is quite typical for this scale (Field et al., 2009). Small-grain species richness is hard to predict, as it depends on so many interacting factors and chance events (Diamond, 1988; Whittaker et al., 2001; Willis and Whittaker, 2002), and small-grained studies typically account for less than 50% of the variation in species richness, even at geographic extents spanning hundreds of kilometers (Field et al., 2009). Not surprisingly, therefore, even our best models left much of the variation unaccounted for, suggesting that other, unmeasured factors also influence woody species richness in the study area. Hydrological, soil factors and biotic interactions might account for some of the residual variation. This is particularly relevant to shrub species richness, which did not correlate strongly with any of our measured variables, and which we could not model well enough to allow prediction. The strongest correlation with shrub species richness was a negative one with basal area, suggesting that shading by trees may reduce shrub diversity. This accords with the recent findings by Oberle et al. (2009) that understorey plant species richness in field plots of similar size to ours correlates much less with regional productivity-related variables than does tree species richness, and that canopy density partly controls shrub species richness at this scale.

Overall, our research contributes to understanding of globally important temperate Andean forests, and represents a step towards targeting conservation of the forests more effectively.

Acknowledgments

This research was made possible by funding from MECESUP project AUS 0103; BIOCORES (Biodiversity conservation, restoration, and sustainable use in fragmented forest landscapes) project ICA-CT-2001-10095; and the FORECOS Scientific Nucleus P04-065-F (MIDEPLAN). We are grateful to Andrés Rivera from Centro de Estudios Científicos (CECS) for the acquisition of the ASTER image needed to develop the research. Thanks to Raúl Bertín and Renato Rivera for assistance with the field work, and Jonathan Barichivich for species identification, all from Universidad Austral of Chile.

Appendix A

List of tree species and shrubs sampled in the study area. Nomenclature follows the Index Kewensis, except for those cases in which no record was found, for which the Gray Herbarium Card Index (http://www.ipni.org) was used.

Species	Family
Trees	
Acacia caven (Molina) Molina	Leguminosae
Aextoxicon punctatum Ruiz & Pav.	Aextoxicaceae
Austrocedrus chilensis (D. Don) Pic.Serm. & M.P.Bizzarri	Cupressaceae
Beilschmiedia berteroana (Gay) Kosterm.	Lauraceae
Crinodendron patagua Molina	Elaeocarpaceae
Cryptocarya alba (Molina) Looser	Lauraceae
Citronella mucronata (Ruiz & Pav.) D.Don	Icacinaceae
Dasyphyllum diacanthoides (Less.) Cabrera	Asteraceae
Drimys winteri J.R. Forst. & G. Forst.	Winteraceae
Embothrium coccineum J.R. Forst. & G. Forst.	Proteaceae
Gevuina avellana Molina	Proteaceae
Kageneckia oblonga Ruiz & Pav.	Rosaceae
Laureliopsis philippiana (Looser) Schodde	Monimiaceae
Laurelia sempervirens (Ruiz & Pav.) Tul.	Monimiaceae
Lithraea caustica Hook. & Arn.	Anacardiaceae
Lomatia dentata R. Br.	Proteaceae
Lomatia hirsuta (Lam.) Diels	Proteaceae
Luma apiculata (DC.) Burret	Myrtaceae
Luma chequen F. Phil.	Myrtaceae

(*continued on next page*)

Author's personal copy

Appendix A (*continued*)

Species	Family
Maytenus boaria Molina	Celastraceae
Nothofagus alpina (Poepp. & Endl.) Oerst.	Fagaceae
Nothofagus dombeyi (Mirb.) Oerst.	Fagaceae
Nothofagus glauca (R. Phil) Krasser	Fagaceae
Nothofagus obliqua (Mirb.) Oerst.	Fagaceae
Nothofagus pumilio (Poepp. & Endl.) Krasser	Fagaceae
Persea lingue (Miers ex Bertero) Nees	Lauraceae
Peumus boldus Molina	Monimiaceae
Quillaja saponaria Molina	Rosaceae
Shrubs	
Acrisione denticulata (Hook. & Arn.) B. Nord.	Asteraceae
Aristotelia chilensis Stuntz	Elaeocarpaceae
Azara celastrina D. Don	Flacourtiaceae
Azara dentata Ruiz & Pav.	Flacourtiaceae
Azara petiolaris (D. Don) I.M.Johnst.	Flacourtiaceae
Azara serrata Ruiz & Pav.	Flacourtiaceae
Baccharis concava Pers.	Asteraceae
Baccharis linearis (Ruiz & Pav.) Pers.	Asteraceae
Baccharis salicifolia (Ruiz & Pav.) Pers.	Asteraceae
Berberis chilensis Gill.	Berberidaceae
Berberis grevilleana Gill.	Berberidaceae
Berberis microphylla G. Forst.	Berberidaceae
Buddleja globosa C. Hope	Buddlejaceae
Cestrum parqui L'Hér.	Solanaceae
Colletia spinosissima J.F. Gmel.	Rhamnaceae
Collihuaja sp 1	Euphorbiaceae
Discaria chacaye (G. Don) Tortosa	Rhamnaceae
Ephedra chilensis C. Presl	Ephedraceae
Undetermined sp1	Escalloniaceae
Fabiana imbricata Ruiz & Pav.	Solanaceae
Gochnatia foliolosa D. Don ex Hook. & Arn.	Asteraceae
Maytenus magellanica Hook.f.	Celastraceae
Mutisia spinosa Hook. & Arn.	Asteraceae
Myoschilos oblongum Ruiz & Pav.	Santalaceae
Myrceugenia ovata O. Berg	Myrtaceae
Pernettia mucronata Gaudich. ex G. Don	Ericaceae
Podanthus mitiqui Lindl.	Asteraceae
Proustia cuneifolia D. Don	Asteraceae
Undetermined sp 2	Rhamnaceae
Ribes cucullatum Hook. & Arn.	Grossulariaceae
Ribes magellanicum Poir.	Grossulariaceae
Schinus montanus Engl.	Anacardiaceae
Schinus patagonicus (Phil.) I.M. Johnst. ex Cabrera	Anacardiaceae
Senna sp 1	Fabaceae
Schinus polygamus (Cav.) Cabrera & I.M. Johnst.	Anacardiaceae
Undetermined sp 3	Solanaceae
Sophora macrocarpa Sm.	Leguminosae
Undetermined sp 4	*Undetermined*
Undetermined sp 5	*Undetermined*

Appendix B. Supplementary material

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.biocon.2010.05.016.

References

Acevedo, D.H., Currie, D., 2003. Does climate determine broad-scale patterns of species richness? A test of the causal link by natural experiment. Global Ecology and Biogeography 12, 461–473.

Algar, A.C., Kharouba, H.M., Young, E.R., Kerr, J.T., 2009. Predicting the future of species diversity: macroecological theory, climate change, and direct tests of alternative forecasting methods. Ecography 32, 22–33.

Altamirano, A., Lara, A., 2010. Deforestación en ecosistemas templados de la precordillera andina del centro-sur de Chile. Bosque 31, 53–64.

Altamirano, A., Echeverría, C., Lara, A., 2007. Efecto de la fragmentación forestal sobre la estructura vegetacional de las poblaciones amenazadas de Legrandia concinna (Myrtaceae) del centro-sur de Chile. Revista Chilena de Historia Natural 80, 27–42.

Anderson, D.R., Burnham, K.P., 1999. Understanding information criteria for selection among capture–recapture or ring recovery models. Bird Study 46, 14–21.

Aplin, P., 2004. Remote sensing: land cover. Progress in Physical Geography 28, 283–293.

Aplin, P., 2005. Remote sensing: ecology. Progress in Physical Geography 29, 104–113.

Araújo, M.B., 1999. Distribution patterns of biodiversity and the design of a representative reserve network in Portugal. Diversity and Distributions 5, 151–163.

Araújo, M.B., New, M., 2007. Ensemble forecasting of species distributions. Trends in Ecology and Evolution 22, 42–47.

Armenteras, D., Rudas, G., Rodriguez, N., Sua, S., Romero, M., 2006. Patterns and causes of deforestation in the Colombian Amazon. Ecological Indicators 6, 353–368.

Armesto, J.J., Rozzi, R., Smith-Ramírez, C., Arroyo, M.T.K., 1998. Conservation targets in South American temperate forests. Science 282, 1271–1272.

Arroyo, M.T.K., Cavieres, L., Peñazola, A., Riveros, M., Faggi, A.M., 1996. Relaciones fitogeográficas y patrones regionales de riqueza de especies en la flora del bosque lluvioso templado de Sudamérica. In: Armesto, J., Villagrán, C., Arroyo, M.T.K. (Eds.), Ecología de los Bosques Nativos de Chile. Editorial Universitaria, Santiago, pp. 71–99.

Austin, M.P., 1980. Searching for a model for use in vegetation analysis. Vegetation 42, 11–21.

Babcock, B.A., Lakshminarayan, P.G., Wu, J.J., Zilberman, D., 1997. Targeting tools for the purchase of environmental amenities. Land Economics 73, 325–339.

Bannari, A., Morin, D., Bonn, F., Huete, A.R., 1995. A review of vegetation indices. Remote Sensing Reviews 13, 95–120.

Bhattarai, K.R., Vetaas, O.R., 2003. Variation in plant species richness of different life forms along a subtropical elevation gradient in the Himalayas, east Nepal. Global Ecology and Biogeography 12, 327–340.

Bini, M., Diniz-Filho, J., Rangel, T., Akre, T., Albaladejo, R., Albuquerque, F., Aparicio, A., Araújo, M., Baselga, A., Beck, J., Bellocq, M., Böhning-Gaese, K., Borges, P., Castro-Parga, I., Chey, V., Chown, S., de Marco, P., Dobkin, D., Ferrer-Castán, D., Field, R., Filloy, J., Fleishman, E., Gómez, J., Hortal, J., Iverson, J., Kerr, J., Kissling, W., Kitching, I., León-Cortés, J., Lobo, J., Montoya, D., Morales-Castilla, I., Moreno, J., Oberdorff, T., Olalla-Tárraga, M., Pausas, J., Qian, H., Rahbek, C., Rodríguez, M., Rueda, M., Ruggiero, A., Sackmann, P., Sanders, N., Terribile, L., Vetaas, O., Hawkins, B., 2009. Coefficient shifts in geographical ecology: an empirical evaluation of spatial and non-spatial regression. Ecography 32, 193–204.

Cayuela, L., Rey Benayas, J.M., Justel, A., Salas, J., 2006a. Modelling tree diversity in a highly fragmented tropical montane landscape. Global Ecology and Biogeography 15, 602–613.

Cayuela, L., Golicher, D.J., Rey Benayas, J.M., González-Espinosa, M., Ramírez-Marcial, N., 2006b. Fragmentation, disturbance and tree diversity conservation in tropical montane forests. Journal of Applied Ecology 43, 1172–1181.

Conaf, Conama, Birf, Universidad Austral de Chile, Pontificia Universidad Católica de Chile, Universidad Católica de Temuco, 1999. Catastro y Evaluación de los Recursos Vegetacionales Nativos de Chile. Informe Nacional con Variables Ambientales. Santiago, Chile.

Insightful Corporation, 2001. S-PLUS 6 for Windows User's Guide. Insightful Corporation, Seattle.

Crawley, M.J., 2002. Statistical Computing. An introduction to data analysis using S-Plus. John Wiley & Sons, Chichester.

Currie, D., 1991. Energy and large-scale patterns of animal and plant species richness. The American Naturalist 137, 27–49.

Currie, D.J., Mittelbach, G.G., Cornell, H.V., Field, R., Guégan, J.-F., Hawkins, B.A., Kaufman, D.M., Kerr, J.T., Ooberdorff, T., O'Brien, E.M., Turner, J.R.G., 2004. Predictions and tests of climate-based hypotheses of broad-scale variation in taxonomic richness. Ecology Letters 7, 1121–1134.

Davis, F., Andelman, S., Stoms, D., 1999. Sites: An Analytical Toolbox for Ecoregional Conservation Planning. The Nature Conservancy. <http://www.biogeog.ucsb.edu/projects/tnc/toolbox.html> (accessed 29.12.09).

De Vries, P.J., Walla, T., Greney, H., 1999. Species diversity in spatial and temporal dimensions of fruit-feeding butterflies from two Ecuadorian rainforests. Biological Journal of the Linnean Society 68, 333–353.

Diamond, J., 1988. Factors controlling species diversity: overview and synthesis. Annals of the Missouri Botanical Garden 75, 117–129.

Dinerstein, E., Olson, D.M., Graham, D.J., Webster, A.L., Primm, S.A., Bookbinder, M.P., 1995. Una evaluación del estado de conservación de las ecoregiones terrestres de América Latina y el Caribe. Banco Mundial/WWF, Washington, DC.

Donoso, C., 1994. Bosques templados de Chile y Argentina. Variación estructura y dinámica. Segunda Edición. Editorial Universitaria, Santiago.

Drechsler, M., Johst, K., Ohl, C., Watzold, F., 2007. Designing cost-effective payments for conservation measures to generate spatiotemporal habitat heterogeneity. Conservation Biology 21, 1475–1486.

Author's personal copy

Echeverría, C., Coomes, D., Salas, J., Rey-Benayas, J.M., Lara, A., Newton, A., 2006. Rapid deforestation and fragmentation of chilean temperate forests. Biological Conservation 130, 481–494.

ERDAS, 1999. Erdas Field Guide, fifth ed. Erdas Inc., Atlanta (revised and expanded).

Field, R., O'Brien, E.M., Whittaker, R.J., 2005. Global models for predicting woody plant richness from climate: development and evaluation. Ecology 86, 2263–2277.

Field, R., Hawkins, B.A., Cornell, H.V., Currie, D.J., Diniz-Filho, J.A.F., Guégan, J.-F., Kaufman, D.M., Kerr, J.T., Mittelbach, G.G., Ooberdorff, T., O'Brien, E.M., Turner, J.R.G., 2009. Spatial species-richness gradients across scales: a meta-analysis. Journal of Biogeography 36, 132–147.

Francis, A.P., Currie, D.J., 2003. A globally consistent richness–climate relationship for angiosperms. The American Naturalist 161, 523–536.

Game, E.T., Grantham, H.S., 2008. Manual del Usuario de Marxan: Para la versión Marxan 1.8.10. Universidad de Queensland, St. Lucia, Queensland, Australia, y la Asociación para la Investigación y Análisis Marino del Pacífico, Vancouver, British Columbia, Canadá. <http://www.uq.edu.au/marxan/> (accessed 29.12.09).

Gao, B.C., 1996. NDWI-A normalized difference water index for remote sensing of vegetation liquid water from space. Remote Sensing of Environment 58, 257–266.

Gonzales, E., Arcese, P., Schulz, R., Bunnell, F., 2003. Strategic reserve design in the central coast of British Columbia: integrating ecological and industrial goals. Canadian Journal of Forest Research 33, 2129–2140.

González-Espinosa, M., Rey-Benayas, J.M., Ramírez-Marcial, N., Huston, M.A., Golicher, D., 2004. Tree diversity in the northern Neotropics: regional patterns in highly diverse Chiapas, Mexico. Ecography 27, 741–756.

Guisan, A., Zimmermann, N.E., 2000. Predictive habitat distribution models in ecology. Ecological Modeling 135, 147–186.

Hall, J., Burgess, N.D., Lovett, J., Mbilinyi, B., Gereau, R.E., 2009. Conservation implications of deforestation across an elevational gradient in the Eastern Arc Mountains, Tanzania. Biological Conservation 142, 2510–2521.

Harrell, Jr. F.E., and with contributions from many other users, 2009. Hmisc: Harrell Miscellaneous. R package version 3.7-0. <http://CRAN.R-project.org/package=Hmisc>.

Hawkins, B.A., Field, R., Cornell, H.V., Currie, D.J., Guégan, J.-F., Kaufman, D.M., Kerr, J.T., Mittelbach, G.G., Ooberdorff, T., O'Brien, E.M., Porter, E.E., Turner, J.R.G., 2003. Energy, water, and broad-scale geographic patterns of species richness. Ecology 84, 3105–3117.

Hechenleitner, P., Gardner, M., Thomas, P., Echeverría, C., Escobar, B., Brownless, P., Martínez, C., 2005. Plantas amenazadas del centro-sur de Chile. Distribución, conservación y propagación. Trama Impresores, Valdivia.

Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G., Jarvis, A., 2005. Very high resolution interpolated climate surfaces for global land areas. International Journal of Climatology 25, 1965–1978.

IPCC, 2001. Climate Change 2001: Synthesis Report (eds. R. Watson, and IPCC core writing team). <http://www.grida.no/climate/ipcc_tar/vol4/english/027.htm>.

Jocqué, M., Field, R., Brendonck, L., de Meester, L., 2010. Climatic control of dispersal–ecological specialization trade-offs: a metacommunity process at the heart of the latitudinal diversity gradient? Global Ecology and Biogeography 19, 244–252.

Lara, A., Donoso, C., Aravena, J.C., 1996. La conservación de los bosques nativos en Chile: problemas y desafíos. In: Armesto, J.J., Villagrán, C., Arroyo, M.K. (Eds.), Ecología de los bosques nativos de Chile. Editorial Universitaria, Santiago, pp. 335–362.

Lara, A., Soto, D., Armesto, J., Donoso, P., Wernli, C., Nahuelhual, L., Squeo, F. (Eds.), 2003. Componentes científicos clave para una política nacional sobre usos, servicios y conservación de los bosques nativos chilenos. Libro resultante de la reunión científica sobre bosques nativos realizada en Valdivia, los días 17–18 julio de 2003. Universidad Austral de Chile. Iniciativa Científica Milenio de Mideplan.

Lawton, J.H., Bignell, D.E., Bolton, B., Bloemers, G.F., Eggleton, P., Hammond, P.M., Hodda, M., Holt, R.D., Larsen, T.B., Mawdsley, N.A., Stork, N.E., Srivastava, D.S., Watt, A.D., 1998. Biodiversity inventories, indicator taxa and effects of habitat modification in tropical forest. Nature 391, 72–76.

Legendre, P., Legendre, L., 1998. Numerical Ecology. Elsevier Science, Amsterdam.

Lillesand, T., Kiefer, R., Chipman, J., 2004. Remote Sensing and Image Interpretation, fifth ed. John Wiley & Sons, New York.

Luoto, M., Toivonen, T., Heikkinen, R., 2002. Prediction of total and rare plant species richness in agricultural landscapes from satellite images and topographic data. Landscape Ecology 17, 195–217.

MacMillan, D.C., Harley, D., Morrison, R.A., 1998. Cost-effectiveness analysis of woodland ecosystem restoration. Ecological Economics 27, 313–324.

Mather, P., 1999. Computer Processing of Remotely-Sensed images. An Introduction, second ed. John Wiley & Sons, Chichester.

May, R.M., Lawton, J.H., Stork, N.E., 1995. Assessing extinction rates. In: Lawton, J.H., May, R.M. (Eds.), Extinction Rates. Oxford University Press, Oxford, pp. 1–24.

Murphy, D.D., 1990. Conservation Biology and scientific method. Conservation Biology 4, 203–204.

Murtaugh, P.A., 2009. Performance of several variable-selection methods applied to real ecological data. Ecology Letters 12, 1061–1068.

Myers, N., Mittermeler, R.A., Mittermeler, C.G., Da Fonseca, G.A., Kent, J., 2000. Biodiversity hotspots for conservation priorities. Nature 403, 853–858.

Nagendra, H., 2001. Using remote sensing to assess biodiversity. International Journal of Remote Sensing 22, 2377–2400.

Newton, A.C., 2007. Biodiversity Loss and Conservation in Fragmented Forest Landscapes. CABI Publishing, Wallingford, Oxford.

Oberle, B., Grace, J.B., Chase, J.M., 2009. Beneath the veil: plant growth form influences the strength of species richness-productivity relationships in forests. Global Ecology and Biogeography 18, 416–425.

O'Brien, E.M., 1998. Water-energy dynamics, climate and prediction of woody plant species richness: an interim general model. Journal of Biogeography 25, 379–398.

O'Brien, E.M., Field, R., Whittaker, R.J., 2000. Climatic gradients in woody plant (tree and shrub) diversity: water energy dynamics, residual variation, and topography. Oikos 89, 588–600.

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Simpson, G.L., Stevens, M.H.H., 2008. Vegan: community ecology package version 1.11-0. <http://cran.r-project.org/> (accessed 17.03.09).

Olivares, P., 1999. Sustitución de bosque nativo por otros usos del suelo en dos sectores de la precordillera andina de la VII región, entre los años 1987 y 1996. Tesis de grado Ingeniería Forestal, Facultad de Ciencias Forestales, Universidad Austral de Chile, Valdivia.

Parolo, G., Rossi, G., 2008. Upward migration of vascular plants following a climate warming trend in the Alps. Basic and Applied Ecology 9, 100–107.

Pauchard, A., Alaback, P., 2004. Influence of elevation, land use and landscape context on patterns of alien plant invasions along roadsides in protected areas of south-central Chile. Conservation Biology 18, 238–248.

Pausas, J.G., Austin, M.P., 2001. Patterns of plant species richness in relation to different environments: an appraisal. Journal of Vegetation Science 12, 153–166.

Peres-Neto, P.R., Legendre, P., Dray, S., Borcard, D., 2006. Variation partitioning of species data matrices: estimation and comparison of fractions. Ecology 87, 2614–2625.

Peterson, A.T., Naksawa, Y., 2008. Environmental data sets matter in ecological niche modelling: an example with *Solenopsis invicta* and *Solenopsis richteri*. Global Ecology and Biogeography 17, 135–144.

Pezoa, L.S., 2003. Recopilación y análisis de la variación de las temperaturas (período 1965–2001) y las precipitaciones (período 1931–2001) a partir de la información de estaciones meteorológicas de Chile entre los 33° y 53° de latitud sur. Tesis de grado Ingeniería Forestal, Facultad de Ciencias Forestales, Universidad Austral de Chile, Valdivia.

Pickering, C., Hill, W., Green, K., 2008. Vascular plant diversity and climate change in the alpine zone of the Snowy Mountains, Australia. Biodiversity Conservation 17, 1627–1644.

Quinn, G., Keough, M., 2003. Experimental Design and Data Analysis for Biologists. Cambridge University Press, Cambridge.

R Development Core Team, 2009. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN: 3-900051-07-0. <http://www.R-project.org>.

Ramírez-Marcial, N., González-Espinosa, M., Williams-Linera, G., 2001. Anthropogenic disturbance and tree diversity in montane rain forests in Chiapas, Mexico. Forest Ecology and Management 154, 311–326.

Rangel, T.F.L.V.B., Diniz-Filho, J.A.F., Bini, L.M., 2006. Towards an integrated computational tool for spatial analysis in macroecology and biogeography. Global Ecology and Biogeography 15, 321–327.

Schlatter, J.E., Gerding, V., Adriazola, J., 1997. Sistema de Ordenamiento de la Tierra: Herramienta para la planificación forestal aplicada a las regiones VII, VIII y IX, second ed. Universidad Austral de Chile, Facultad de Ciencias Forestales, Valdivia.

Smith-Ramírez, C., 2004. The Chilean coastal range: a vanishing center of biodiversity and endemism in South American temperate rainforests. Biodiversity and Conservation 13, 373–393.

Srivastava, D.S., Lawton, J.H., 1998. Why more productive sites have more species: an experimental test of theory using treehole communities. The American Naturalist 152, 510–529.

Trivedi, M., Morecroft, M.D., Berrya, P.M., Dawson, T.P., 2008. Potential effects of climate change on plant communities in three montane nature reserves in Scotland, UK. Biological Conservation 141, 1665–1675.

Turner, W., Spector, S., Gardiner, N., Fladeland, M., Sterling, E., Steininger, M., 2003. Remote sensing for biodiversity science and conservation. Trends in Ecology and Evolution 18, 306–314.

Van der Horst, D., 2007. Assessing the efficiency gains of improved spatial targeting of policy interventions: the example of an agri-environmental scheme. Journal of Environmental Management 85, 1076–1087.

Venables, W.N., Ripley, B.D., 2002. Modern Applied Statistics with S. Springer-Verlag, New York.

Venevsky, S., Veneskaia, I., 2003. Large-scale energetic and landscape factors of vegetation diversity. Ecology Letters 6, 1004–1016.

Whittaker, R.J., Willis, K.J., Field, R., 2001. Scale and species richness: towards a general, hierarchical theory of species diversity. Journal of Biogeography 28, 453–470.

Willis, K.J., Whittaker, R.J., 2002. Species diversity – scale matters. Science 295, 1245–1248.

Zhang, X.C., Nearing, M.A., 2005. Impact of climate change on soil erosion, runoff, and wheat productivity in central Oklahoma. Catena 61, 185–195.